



The Application of Response Surface Methodology (RSM) In the Computational Optimization of Sustained Release (SR) For Phenothiazine Derivative Matrix Tablet

Shiva Kant Thakur ^{a*}, Rahul Pal ^a, Devanand Jha ^{b++},
Prottay Dutta ^{c#}, Prachi Pandey ^a,
Bhuneshwar Dutta Tripathi ^{b†}, Mohammad Rizwan ^{d#},
Shradha Ojha ^{e++}, Divya ^{e++} and Ravindra Pal Singh ^a

^a Department of Pharmaceutics, Nims Institute of Pharmacy, Nims University Rajasthan, Jaipur, 303121, India.

^b Department of Pharmacology, Narayan Institute of Pharmacy, Gopal Narayan Singh University, Jamuhar, Rohtas, Bihar-821305, India.

^c Department of Pharmacy, Usha Martin University, Ranchi, Jharkhand-835103, India.

^d Department of Pharmaceutical Science, Sir J.C. Bose Technical Campus, Bhimtal, Kumaun University, Nainital-263136, India.

^e Invertis Institute of Pharmacy, Invertis University, Bareilly, U.P., India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2023/v35i357483

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111654>

Original Research Article

Received: 23/10/2023
Accepted: 28/12/2023
Published: 28/12/2023

⁺⁺ Research Scholar;

[#] Assistant Professor;

[†] Associate Professor;

*Corresponding author: E-mail: palsrahul330@gmail.com;

ABSTRACT

Introduction: This study explores the use of Response Surface Methodology (RSM), a statistical optimization technique, to optimize the SR properties of prochlorperazine maleate (PCM) matrix tablets. PCM is a phenothiazine derivative used for treating schizophrenia, nausea, and vomiting. Sustained-release formulations offer extended drug delivery, potentially improving patient compliance and reducing side effects. RSM helps identify optimal combinations of critical formulation factors influencing drug release, such as polymer type and concentration, filler type, and drug/polymer ratio. The study likely involves designing experiments based on chosen RSM designs (e.g., Box-Behnken) with varying factor levels. Formulate SR tablets with different factor combinations. Evaluating the drug release profiles of each tablet formulation. Analyzing data using RSM software to build mathematical models relating factors to drug release and identifying optimal factor combinations that maximize desired release characteristics.

Objective: The ongoing research purpose to improve the advancement of a sustained release tablet containing Phenothiazine derivative PCM loaded matrix. This is achieved by utilizing DoE as a computational method to statistically validate the formulation.

Methodology: Basically two methods involved in the formulation using direct compression; a simple and efficient method where all dry ingredients are blended and directly compressed into tablets.

Results: The obtained outcomes closely matched the anticipated values derived from the experimental setup. The enhanced PCM matrix tablets demonstrated a prolonged and uniform discharge of PCM over a span of 6 hours.

Conclusion: This study Based on a 2³ FD, response surface methodology was used to successfully develop the current research of PCM matrix tablets for sustained release application. The corresponding contour plots and 3D response methodology represent the important variables for the optimisation process.

Keywords: Sustained release; tablets; statistical analysis; contour plot; response surface methodology; optimization; antiemetic formulations; computational approach; factorial study; design of experiments; and designs.

1. INTRODUCTION


The matrix tablet formulated for sustained release refers to a specific type of medication where the active ingredient is dispersed throughout a solid, inert matrix. This matrix acts as a controlled-release mechanism, gradually releasing the drug over a longer period of time compared to a conventional immediate-release tablet [1-2]. The mechanism is the drug is that physically dispersed within the matrix, usually composed of polymers and other excipients. The matrix allows for slow diffusion of the drug into the surrounding fluids, controlling the release rate.

RSM is a powerful statistical and mathematical tool widely employed in the pharmaceutical industry, particularly for optimizing the formulation of matrix tablets. It helps to identify the optimal combination of critical factors influencing the desired characteristics of the tablet, such as drug release rate, hardness, and disintegration time [3-4]. DoE plays a crucial role in both formulation development and optimization. It helps researchers systematically

explore the influence of various factors on desired responses, paving the way for efficient and effective formulation design.

PCM, also known by the brand name Compazine, is a medication belonging to the phenothiazine class. It acts primarily as an antiemetic (prevents nausea and vomiting) and an antipsychotic (treats mental disorders like schizophrenia) [5]. The chemical structure of PCM shown in the given Fig. 01 as below followings:

Phenothiazines are a group of older antipsychotic medications with various applications. They share a similar chemical structure and mechanism of action, but differ in potency and side effect profiles. PCM is considered a first-generation phenothiazine, meaning it was developed earlier and has a higher risk of side effects compared to newer generations [6-7]. Overall, PCM is an effective medication for managing nausea and vomiting and short-term treatment of certain mental disorders. However, it's crucial to be aware of its potential side effects and use it only under the guidance of a healthcare professional [8].



**2-chloro-10-[3-(4-methylpiperazin-1-yl)propyl]phenothiazine
maleate**

Fig. 1. The chemical structure of prochlorperazine maleate (PCM) with its IUPAC name

Table 1. The PCM matrix tablet formulation batches for each suggested trial formulation

Preparation Batches	API (mg)	Calcium alginate (mg) (M)	EC (mg) (N)	Carbopol 971 (mg) (R)	Lactose (mg)	Mg-Stearate (mg)
RP1	25	16 (+1)	0 (-1)	0 (-1)	85	10.5
RP2	25	32 (-1)	25 (+1)	25 (+1)	85	10.5
RP3	25	32 (-1)	0 (-1)	0 (-1)	85	10.5
RP4	25	32 (+1)	25 (+1)	25 (-1)	85	10.5
RP5	25	16 (+1)	25 (+1)	25 (+1)	85	10.5
RP6	25	16 (-1)	0 (-1)	25 (+1)	85	10.5
RP7	25	32 (+1)	0 (-1)	25 (+1)	85	10.5
RP8	25	16 (1)	25 (+1)	0 (1)	85	10.5

Where; EC: Ethyl cellulose.

Manufacturing a pharmaceutical formulation that delivers ideal quality quickly and with few trials is essential for creating a matrix tablet for continuous release, for example. Due to its ability to reduce the necessity for significant research, the RSM is a generally used strategy for creating and optimising diverse pharmacological formulations [8-10].

The primary objective of the ongoing research is to enhance the progress of a sustained release tablet that contains a PCM loaded matrix. This goal is accomplished by employing DoE as a computational method to statistically validate the formulation. An investigation was conducted using a computer-aided optimization technique, specifically a 2^3 FD, to analyze the impact of different amounts of hydrophilic polymers on the properties of PCM sustained release matrix tablets [10]. The three independent process variables, namely calcium alginate, EC, and Carbopol 971, were considered as factors in the polymer-blend. The study focused on evaluating the effects of these factors on drug release and tablet hardness [11-12].

2. METHODOLOGY / MATERIALS

2.1 Chemical Agents

Lactose and prochlorperazine maleate (PCM) were acquired from Pvt. Ltd. in India. We bought carbopol 971, ethyl cellulose (EC), and calcium alginate from M.K. Specialties Pvt. Ltd. in India. Analytical grade reagents and chemicals were utilised in all other instances.

2.2 Formulation of PMC loaded Matrix Tablet

PCM matrix tablets were prepared using the direct compression approach. In order to do this, different excipients were combined with hydrophilic polymers in the proper ratios to act as release modifiers. Using an #80 sieve, the medication, polymers, and excipients were filtered. After that, a single punch tablet machine from Pvt. Ltd., India was used to properly blend and compress the medicine together with all the other elements. For a batch size of 100 tablets, 6

mm round and flat punches were used to compress the material.

2.3 Experimental Design (2³ Factorial Design)

The FD with three factors and two levels was employed to optimize the PCM matrix tablet. The independent variables, namely the amount of calcium alginate (M), EC (N), and Carbopol 971 (R) in the polymer-blend, were varied at low and high levels. Several trial formulations of the PCM matrix tablets were created based on the FD's trial suggestion. The formulation chart for each suggested trial is shown in the table. The hardness (kg/cm²) and cumulative drug release after 6 hours (R6h. %) were examined as dependent variables. Design-Expert 8.0.6.1 was utilized to construct and evaluate the statistical experimental design [13-15]. This design matrix, which includes the investigated factors and responses, is shown in Table. 02.

2.4 Determination of Drug Content

Twenty tablets from each batch of formulation were weighed and crushed to a fine powder. The powder equivalent to 20 mg of PCM was then transferred to a 100 mL volumetric flask. In order to ensure complete dissolution, the flask was filled up to the 100 mL mark with 0.1 N HCl, while vigorously shaking the contents. The resulting mixture was filtered using filter paper no. 40. Following appropriate dilution, the absorbance values were measured at the maximum λ max of 254.5 nm using a UV-VIS spec [16-17].

2.5 Weight Variation Test

From each batch, twenty tablets were selected, and an electronic analytical balance was used to measure the tablets precisely. Next, the percentage of weight fluctuation was computed.

$$\text{Weight variation (\%)} = \frac{\text{Standard Deviation}}{\text{Mean Weight}} \times 100$$

2.6 Hardness Test

The PCM matrix tablets' hardness was assessed using the Pfizer hardness tester. Initially, the tester was adjusted to zero and the tablets were placed between two jaws. The tablet was then forced to shatter into smaller pieces. The tester's scale was examined to determine the kilogram-level pressure needed to break the pill [18].

2.7 In-Vitro Drug Release (DR) Evaluation

The *in-vitro* DR studies were conducted using the USP dissolution apparatus, specifically the

basket type manufactured by Campbell Electronics in India. The experiments were performed at a speed of 50 rpm and a temperature of $37 \pm 0.5^\circ\text{C}$. PCM matrix tablets were placed in the basket and the dissolution process involved using 900 mL of 0.1 N HCl (pH 1.2) for the initial 2 hours, followed by 900 mL of phosphate buffer (pH 7.4) for the remaining duration. To keep track of the PCM release, 5 mL portions were taken out from the disso. Container at specific time intervals and replaced with the same amount of new solution. The gathered samples were then filtered using Whatman filter paper (no. 40) and analyzed for PCM content using a UV-VIS spec. made by Thermo Spectronic in the USA, at a wavelength of 254.5 nm [19-21].

2.8 Kinetic Analysis of Release Data

In order to examine how drugs are released from PCM matrix tablets, the *in vitro* dissolution data was analyzed using different mathematical models [22-23]. These models include the zero-order, first-order, Higuchi, and Korsmeyer-Peppas models as in Table. 03 as below followings:

The Korsmeyer-Peppas model has once again been used to analyze drug release behavior in different pharmaceutical formulations. Its purpose is to distinguish between various releases mechanisms like Fickian release, non-Fickian release, and case-II transport. Fickian release occurs when the value of *n* is 0.5 or less [24]. Non-Fickian release is defined when the value of *n* is between 0.4 and 1.0. On the other hand, case-II transport happens when the value of *n* is 1.0 or greater, involving polymer dissolution and polymeric chain enlargement or relaxation.

2.9 Statistical Analysis (SA)

The SA is the science of collecting, examining, and interpreting data to uncover patterns, trends, and relationships. It's a powerful tool used in various fields, from scientific research and social sciences to business and healthcare. The statistical optimization was conducted utilizing Design-Expert 8.0.6.1 software developed by Stat-Ease Inc. Simple statistics were employed to analyze all other data [24-26].

2.10 Optimization of Receptor and Ligands for PCM

A phenothiazine derivative belonging to the class of atypical antipsychotics. Primarily used to treat

nausea, vomiting, and vertigo associated with motion sickness and chemotherapy. Also prescribed for schizophrenia, anxiety, and other psychiatric disorders.

2.11 PCM and its Receptor Binding

PCM interacts with various receptors in the central nervous system, including dopamine D2, serotonin 5-HT2, and histamine H1 receptors.

The antiemetic effects are mainly attributed to dopamine D2 receptor antagonism in the chemoreceptor trigger zone (CTZ). Antagonism of serotonin 5-HT2 receptors contributes to its antipsychotic and anxiolytic properties [26-27].

2.12 Optimization of Receptor and Ligands for PCM

Receptor optimization aims to improve the binding affinity and selectivity of PCM towards its target receptors. This can be achieved through:

- **Structure-based drug design:** Utilizing computational modeling and structural information of the receptors to design PCM analogs with enhanced binding properties.
- **High-throughput screening:** Identifying new ligands from large compound libraries that bind more potently and selectively to the desired receptors [28].

2.13 Ligand optimization Focuses on Modifying the Chemical Structure of PCM to

- Increase its binding affinity to the target receptors.
- Improve its selectivity over other receptors, minimizing side effects.
- Enhance its pharmacokinetic properties, such as oral bioavailability and blood-brain barrier penetration.

3. RESULTS AND DISCUSSION

3.1 Optimization of PCM Matrix Tablet

Pharmaceutical formulators traditionally create different formulations by changing one factor at a time, which is a time-consuming method. However, this approach often leads to unsuccessful experiments due to poor planning and design. A lack of appropriate planning cannot be compensated for, not even with extensive data analysis. Consequently, it is

critical to comprehend how formulation parameters impact the calibre of formulations that require few experimental trials. As a result, the selection of formulation variables should be based on established statistical tools for optimization. In the case of PCM matrix tablets, a total of 8 trial formulations were proposed using a 2³ FD. These formulations involved three independent variables: the amounts of calcium alginate (M), EC (N), and Carbopol 971 (R), considered in mg quantity, which were varied at two different levels (high and low) [27-29].

The purpose of this study was to analyze how independent variables affect the release of drugs after 6 hours, as well as the percentage and hardness of the tablets. These parameters were considered important for optimization. The used of a 2³ FD to create different trial formulations of PCM matrix tablets using the direct compression method [30]. These ingredients used are listed in Table. 02. The results of the experiments and the corresponding responses are summarized in Table. 03. The ANOVA results confirmed the significance of these models for all response parameters, as shown in Table. 04.

The Design-Expert 8.0.6.1 software used the data to create polynomial model equations that included main factors and interaction factors. The equation showing the relationship between R and 6h% can be rewritten in the following ways:

$$R \text{ 6h (\%)} = 85.36 - 0.66G - 0.85H - 0.79J + 6.6 \times 10^{-3} GH + 1.3 \times 10^{-2} GJ + 9.4 \times 10^{-3} HJ, \\ R^2 = 0.9999; F \text{ value} = 2815.99; P < 0.05 \\ \text{Equation} \quad \quad \quad [3]$$

The model equation, which describes the connection between the response variable hardness, has now taken a different form as followings:

$$\text{Hardness kg/cm}^2 = 3.86 + 0.03G + 8.86 \times 10^{-3} H + 0.06J + 2.49 \times 10^{-4} GH - 1.09 \times 10^{-3} GJ - 1.16 \times 10^{-4} HJ$$

$$R^2 = 0.9997; F \text{ value} = 456.19; P < 0.05 \\ \text{Equation} \quad \quad \quad [4]$$

A simplified model was produced by eliminating non-significant components ($P > 0.05$) from multiple regression analysis-derived model equations during the model simplification phase [29-31].

$$R6h. (\%) = 85.39 - 0.69G - 0.89H - 0.78J + 9.5 \times 10^{-3} HJ, \text{ Equation [5]}$$

$$\text{Hardness } \text{kg/cm}^2 = 2.89 + 0.04G + 7.89 \times 10^{-3} H + 0.08 J - 1.09 \times 10^{-3} GJ$$

The Fig. 03 and Fig. 04 present Pareto charts that were used to analyze the statistical significance of each response coefficient (R6h % and Hardness) respectively.

These charts show the importance of each coefficient visually. Coefficients with t values above the Bonferroni line are considered important, while coefficients with t values between the Bonferroni line and the t limit line are likely to be important. However, coefficients with t values below the t limit line are not statistically significant. These Pareto charts also

help confirm the ANOVA resulting by identifying and removing terms that are not significant ($P > 0.05$) in the model equations [32].

RSM is a very successful method that is applied to the creation and improvement of drug delivery systems. The 3D RSP and the accompanying CP are shown in Fig. 05, 06, 07, 08, 09, and 10, and they aid in estimating the impacts of independent variables on each response under investigation.

These plots are valuable tools for understanding the main and interaction effects of the variables. Additionally, the 2D CP provides a visual representation of the response values. In the case of hardness, as shown in Fig. 03, the Pareto chart reveals both positive and negative effects [30-31].

Table 2. The observed response values with drug contents in PCM matrix tablets and the 2³ factorial designs (FD) are presented

Formulation Batches	Calcium alginate (mg) <i>M</i>	EC (mg) <i>N</i>	Carbopol 971 (mg) <i>R</i>	Observations	
				R 6h (%) ^x	Hardness (kg/cm ²) ^y
RP1	16 (-1)	0 (-1)	0 (-1)	75.72 ± 3.37	4.20 ± 0.26
RP2	32 (+1)	25 (+1)	25 (+1)	57.25 ± 2.14	3.19 ± 0.16
RP3	32 (-1)	0 (-1)	0 (-1)	74.17 ± 2.55	4.53 ± 0.22
RP4	32 (+1)	25 (+1)	0 (-1)	62.42 ± 3.26	3.85 ± 0.17
RP5	16 (-1)	25 (+1)	25 (+1)	51.49 ± 2.18	4.12 ± 0.12
RP6	16 (+1)	0 (+1)	25 (-1)	63.44 ± 1.83	3.89 ± 0.35
RP7	32 (-1)	0 (-1)	25 (+1)	56.86 ± 2.33	4.92 ± 0.14
RP8	16 (-1)	25 (-1)	0 (+1)	60.16 ± 1.74	3.42 ± 0.26

The cumulative drug release after 6 hours is denoted as x R 6h (%). The mean value, represented by y, is accompanied by the standard deviation (S.D.) and is based on a sample size of 6 (n = 6). The main effects (factors) are represented by A, B, and C. A higher value is indicated by (+1), while a lower value is indicated by (-1)




Fig. 2. The Structure of the dopamine D2 receptor in complex with the PCM drug [A]. Chemical structure of PCM and [B]. The docking of PCM

Table 3. The various models with their formula, description and various applications

Model	Description	Formula	Application
Zero-Order	Constant drug release rate, independent of drug concentration	$Q_t = k_0t$	Matrix tablets with low soluble drugs, osmotic systems, transdermal systems
First-Order	Release rate proportional to remaining drug concentration	$\log(Q_t) = kt/2.303$	Water-soluble drugs in porous matrices
Higuchi	Release rate controlled by diffusion through a dissolving matrix	$Q_t = kH\sqrt{t}$	Matrix tablets, coated pellets, transdermal patches
Korsmeyer-Peppas	Release mechanism involves both diffusion and erosion	$M_t/M_\infty = kt^n$	Polymeric systems, swelling devices, capsules

Table 4. The list of statistics of ANOVA generated response for R6h and hardness

Derivation	Composition of R ²	d.f. ^x	Mean R ²	F value	P value prob. > F
<i>R_{6h} (%)</i> ^y					
Models	594.66	6	99.95	2516.98	0.0249 (K)
X	99.85	1	97.85	2422.75	0.0124 (K)
Y	334.08	1	324.08	8764.86	0.0069 (K)
Z	155.19	1	156.19	4214.12	0.0094 (K)
XY	2.95	1	1.93	54.25	0.0763 (NK)
XZ	6.59	1	5.49	152.09	0.0416 (NK)
YZ	6.16	1	6.24	187.02	0.0452 (K)
<i>Hardness (kg/cm²)</i>					
Models	0.87	6	0.16	455.16	0.0451 (K)
X	0.12	1	0.12	403.77	0.0466 (K)
Y	0.15	1	0.14	394.05	0.0425 (K)
Z	0.59	1	0.69	1950.05	0.0159 (K)
XY	2.82×10^{-3}	1	2.82×10^{-3}	8.00	0.3149 (NS)
XZ	0.06	1	0.07	170.03	0.0588 (S)
YZ	1.04×10^{-3}	1	1.04×10^{-3}	4.25	0.4429 (NS)




Fig. 3. The schematic representation of design expert software for R6h. %

**Design-Expert software
hardness**




Fig. 4. The schematic representation of design expert software for hardness




Fig. 5. The RSP [A] and CP [B] illustrate the impact of varying quantities of calcium alginate and EC on the R6h (%) outcome

More calcium alginate, EC, and carbopol 971 in PCM matrix tablets results in lower R 6h % values and higher hardness values, as seen by the 3D RSP and CP for R6h and hardness. To

created optimised PCM matrix tablets using the direct compression approach in order to evaluate the optimisation capabilities of the models created based on a 2^3 FD [32]. One of the

carefully chosen ideal process variable configurations suggested by the experimental design was employed.

The optimized formulation of PCM matrix tablets involved using M = 16.29 mg, N = 34.11 mg, and R = 30.32 mg as selected optimal process variable settings. To determine the optimal values of responses based on the desirability

criterion, we conducted numerical analysis with the assistance of Design expert 8.0.6.1 software [33]. This analysis led to the development of optimized PCM matrix tablet.

The actual values observed during the practical experiments are presented in Table. 05 along with the projected values derived from the mathematical model [33-34].




Fig. 6. The RSP [A] and CP [B] illustrate the impact of varying quantities of calcium alginate and Carbopol 971 on R6h (%)




Fig. 7. The RSP [A] and CP [B] illustrate the impact of varying quantities of EC and Carbopol 971 on R6h (%)




Fig. 8. The RSP [A] and CP [B] illustrate the impact of varying amounts of calcium alginate and EC on the hardness (kg/cm²)




Fig. 9. The RSP [A] and CP [B] show how different concentrations of Carbopol 971 and calcium alginate affect hardness (kg/cm²)

Table 5. The observations obtained from experiments to confirm optimization aptitude

Formulation Code	Calcium alginate (mg) [M]	EC (mg) [N]	Carbopol 971 (mg) [R]	Observations	
				R_{6h} (%) ^x	Hardness (kg/cm ²)
RO	16.29	33.16	31.34	Actual values ^y	
				42.62 ± 1.89	4.76 ± 0.08
				Predicted values	
				39.98	5.59
				% Error	4.04 2.43

*a*R_{6h} (%): DR accumulated after 6 hours; *b* mean ± S.D. of actual values, *n* = 3; *c*% error = (actual value - predicted value)/predicted value × 100; A, B, and C denote the primary factors.




Fig. 10. The RSP [A] and contour plot (CP) [B] illustrate the impact of varying quantities of EC and Carbopol 971 on the hardness (kg/cm²)

The PCM matrix tablets that were optimized showed an R 6h of $42.64 \pm 1.96\%$ and a hardness of $4.67 \pm 0.08 \text{ kg/cm}^2$. The error values were small, less than 5, which suggests that the mathematical models derived from the 2^3 FD were a good fit [35].

0.07 to $4.08 \pm 0.29\%$, with none of them deviating by more than 5% from their average weight (Table. 06) [34-35]. The uniform mixing of CPM with other ingredients is evident from the consistent drug content and weight of the formulated PCM matrix tablets.

3.2 Optimization of Drug Content and Weight Variation

All matrix tablets containing PCM were analyzed and found to contain PCM within the range of 97.99 ± 0.85 to 99.03 ± 0.59 (Table. 06). This confirms that the drug content is present in the appropriate amount in all the matrix tablets. The weight variation of the tablets also indicates adherence to GMPs by the manufacturers and the amount of APIs in the formulation. This PCM loaded matrix tablets meet the weight uniformity specifications outlined in the USP. The tablets exhibited a weight variation ranging from $2.79 \pm$

3.2.1 Hardness

The purpose of conducting the hardness test on PCM matrix tablets was to evaluate their ability to endure handling without breaking or chipping. It has been determined that a minimum force of approximately 4 kg/cm^2 is necessary for the tablets to possess satisfactory hardness. The results obtained from the hardness evaluation indicated that the hardness of the matrix tablets was found to be within the acceptable range of 3.20 ± 0.26 and $5.67 \pm 0.08 \text{ kg/cm}^2$ (Table. 03 and 04) [35-36]. These findings confirm that the tablets possess the desired level of hardness.

Table 6. The Drug content and weight variation of PCM matrix tablets

Formulation codes	Drug content (%) ^x	Weight variation (%) ^y
RP1	98.04 ± 0.61	2.14 ± 0.19
RP2	97.43 ± 0.71	3.36 ± 0.32
RP3	98.25 ± 0.66	2.15 ± 0.31
RP4	99.98 ± 0.84	3.06 ± 0.32
RP5	99.22 ± 1.24	3.42 ± 0.16
RP6	97.64 ± 0.54	2.89 ± 0.16
RP7	97.81 ± 0.77	2.04 ± 0.15
RP8	99.63 ± 0.56	3.84 ± 0.15
RO	97.84 ± 0.73	1.69 ± 0.09

Where; x Mean \pm S.D., n = 20; y: coefficient of weight variation (%) [The weight variation coefficient (%) can be calculated by dividing the standard deviation by the mean weight and then multiplying the result by 100]




Fig. 11. The drug release from different PCM matrix tablets (RP1 to RO) was evaluated in-vitro. The values are expressed as the mean \pm standard deviation (SD), with a sample size

Table 7. The outcomes of curve fitting for the release data of *In-vitro* PCM from various PCM matrix tablets

Formulation Batches	Correlation coefficient (R^2)				
	Zero Order	First Order	Higuchi	Korsmeyer-Peppas	Release exponent (n)
RP1	0.9645	0.9807	0.8278	0.9510	0.61
RP2	0.9798	0.9930	0.7069	0.9416	0.74
RP3	0.9577	0.9952	0.6844	0.8988	0.70
RP4	0.9810	0.9949	0.7288	0.9418	0.71
RP5	0.9733	0.9942	0.7062	0.9084	0.71
RP6	0.9677	0.9908	0.6925	0.9048	0.72
RP7	0.9774	0.9954	0.7059	0.9093	0.72
RP8	0.9670	0.9944	0.7214	0.9098	0.69
RO	0.9744	0.9920	0.7211	0.9470	0.75

3.2.2 In-Vitro Drug Release (DR)

All PCM matrix tablets exhibited drug release in 0.1 N HCl (pH 1.2) for the initial 2 hours, followed by release in phosphate buffer (pH 7.4) for 4 hours. These matrix tablets consistently demonstrated prolonged and sustained drug release for a total of 6 hours (Fig. 11). The cumulative DR from these tablets after 6 hours of dissolution (R6h) ranged from 41.61 ± 1.97 to $74.62 \pm 2.36\%$. Analysis of the response surface revealed that the R6h% values decreased as the three independent variables (amount of calcium alginate, EC, and Carbopol 971) increased Table. 07. The presence of larger quantities of

hydrophilic polymers, such as calcium alginate, EC, and Carbopol 971, leads to an elevated viscosity [37]. This increased viscosity can promote the creation of extremely thick gels when exposed to the aqueous fluids in the dissolution medium. Consequently, this process would decelerate the rate at which drugs are released from the PCM matrix tablets. The release exponent (n) value, obtained from in-vitro PCM release data for different matrix tablets, varied between 0.49 and 0.76. This range suggests the presence of an anomalous diffusion mechanism for drug release. The anomalous diffusion mechanism indicates a combination of

diffusion-controlled and swelling-controlled DR from PCM matrix tablets [38-41].

4. CONCLUSION

The SR application of PCM matrix tablets was successfully achieved through the use of RSM based on a 23-FD. The analysis and optimization of the properties of the matrix tablet, including drug release and hardness, were conducted in a comprehensive manner, focusing on the effects of calcium alginate, EC, and Carbopol 971. Through the examination of RSP and CP, it was observed that higher quantities of calcium alginate, EC, and Carbopol 971 led to reduced drug release (R 6h%) values and increased hardness values. The PCM matrix tablets (RO) that were optimized displayed a drug release rate of $44.72 \pm 1.89\%$ within a span of 6 hours, along with a hardness value 4.86 ± 0.08 kg/cm². This final optimized tablet showcased an extended sustained release of PCM for a period of 12 hours, which has the potential to enhance patient adherence by decreasing the frequency of dosing when compared to traditional tablets.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Zalte HD, Saudagar RB. Review on sustained release matrix tablet. *International Journal of Pharmacy and Biological Sciences*. 2013;3(4): 17-29.
2. Ganesh GNK, Sureshkumar R, Jawahar N, Senthil V, Venkatesh DN, Srinivas MS. Preparation and evaluation of sustained release matrix tablet of diclofenac sodium using natural polymer. *Journal of pharmaceutical sciences and research*. 2010;2(6):360.
3. Alhalmi A, Altowairi M, Saeed O, Alzubaidi N, Almoiliqy M, Abdulmalik W. Sustained release matrix system: an overview. *World J Pharm Pharm Sci*. 2018;7(6):1470-86.
4. Prakhar A, Akhtar S. A comprehensive review on sustained release matrix tablets: a promising dosage form. *Universal Journal of Pharmaceutical Research*. 2018;3(6):49-54.
5. Nautyal U, Gupta D. Oral sustained release tablets: an overview with a special emphasis on matrix tablet. *International Journal of Health and Biological Sciences*. 2020;3(1):06-13.
6. Khuri AI, Mukhopadhyay S. Response surface methodology. *Wiley Interdisciplinary Reviews: Computational Statistics*. 2010;2(2):128-149.
7. Pal R, Pandey P, Rizwan M, Koli M, Thakur SK, Malakar RK, Chawra HS. The Utilization of Response Surface Methodology (RSM) In the Optimization of Diclofenac Sodium (DS) Liposomes Formulate through the Thin Film Hydration (TFH) Technique with Involving Computational Method. *Journal of Advances in Medicine and Medical Research*. 2023;35(22): 287-300.
8. Politis N, S, Colombo P, Colombo G, M. Rekkas D. Design of experiments (DoE) in pharmaceutical development. *Drug development and industrial pharmacy*. 2017;43(6):889-901.
9. Albadry AA, Ali WK, Al-saady FA. Formulation and evaluation of prochlorperazine maleate sustained release floating tablet. *Int J Pharm Pharm Sci*. 2017;9(2):89-98.
10. Pal R, Pandey P, Jha D, Dutta P, Sahoo S, Gupta R, Chawra HS. The Utilization of 32Full Factorial Design (FFD) for Optimization of Lincomycin Hydrochloride (LNH) Loaded Nanogel Involving; Design of Experiments (DoE) an Advanced Approach. *Advances in Research*. 2023; 24(6):272-281.
11. Vaja Divyeshkumar N, Patel Maulik M, Joshi Ujjwal T, Patel Jaykishan M. Design, development and characterization of orally disintegrating tablet of prochlorperazine maleate. *J. Chem*. 2010; 2(5):307-312.
12. Malakar J, Nayak AK. Formulation and statistical optimization of multiple-unit ibuprofen-loaded buoyant system using 23-factorial design. *Chemical Engineering Research and Design*. 2012; 90(11):1834-1846.
13. Pandey P, Pal R. Bharath KS, Akilandeshwari A, Koli M, Rizwan M, Chawra HS. Design of Experiments (DoE) manipulation in the formulation and optimization of a traditional Ayurvedic

- medicine derived from dried extract of Senegalia catechu enhanced through statistical analysis. Journal of Pharmacognosy and Phytochemistry. 2023;12(6):159-169.
14. Savaşer A, Özkan Y, Işimer A. Preparation and in vitro evaluation of sustained release tablet formulations of diclofenac sodium. *Il Farmaco*. 2005; 60(2):171-177.
 15. Ofori-Kwakye K, Osei-Yeboah F, Kipo SL. Formulation and quality evaluation of two conventional release tablet formulations. *International Journal of Pharmaceutical Sciences Review and Research*. 2010;4(1):94-99.
 16. Pandey, Prachi & Pal, Rahul & Thakur, Shiva Kant & Sharma, Vidhi & Chanana, Arsh & Singh, Ravinder. future aspects & modification in carbon- nanoparticles in treatment and diagnosis. *World journal of pharmacy and pharmaceutical sciences*. 2023;12:307-324. DOI:10.20959/wjpps20232-24094.
 17. Bohrey S, Chourasiya V, Pandey A. Polymeric nanoparticles containing diazepam: preparation, optimization, characterization, in-vitro drug release and release kinetic study. *Nano Convergence*. 2016;3(1):1-7.
 18. Clark AL, Johnson M, Fairhurst C, Torgerson D, Cockayne S, Rodgers S, Greenstone M. Statistical analysis. In Does home oxygen therapy (HOT) in addition to standard care reduce disease severity and improve symptoms in people with chronic heart failure? A randomised trial of home oxygen therapy for patients with chronic heart failure. *NIHR Journals Library*; 2015.
 19. Pal R, Pandey P, Nogai L, Anand A, Suthar P, SahdevKesar M, Kumar V. The future perspectives and novel approach on gastro retentive drug delivery system (GRDDS) with currrent state. *Journal of Population Therapeutics and Clinical Pharmacology*. 2023;30(17): 594-613.
 20. Kenakin T. Predicting therapeutic value in the lead optimization phase of drug discovery. *Nature Reviews Drug Discovery*. 2003;2(6):429-438.
 21. Pal R, Pandey P, Nogai L. The Advanced Approach in The Development of Targeted Drug Delivery (TDD) With Their Bio-Medical Applications: A Descriptive Review. *International Neuroulogy Journal*. 2023;27(4):40-58.
 22. Wermuth CG. Selective optimization of side activities: another way for drug discovery. *Journal of medicinal chemistry*. 2004;47(6):1303-1314.
 23. Caldwell GW, Yan Z. Optimization in drug discovery. New York: Springer.[Google Scholar]; 2014.
 24. Pal, Rahul Pandey, Prachi Kumar Maurya, Vishesh & Saxena, Archita & Koli, Manju. Optimization And Formulation of Doxorubicin (DOX) Loaded Liposome Well-Used in Chemotherapy Involving Quality by Design (QBD): A Transitory Research. *European Chemical Bulletin*. 2023;12:4491-4510. DOI:10.48047/ecb/2023.12.si5a.0355.
 25. Dua'a KA, Kamal BA. Formulation and Optimization of Oral Fast Dissolving Prochlorperazine Maleate Tablets. *Iraqi Journal of Pharmaceutical Sciences*; 2012. (P-ISSN 1683-3597 E-ISSN 2521-3512), 21(1), 46-55.
 26. Pal, Rahul & Pandey, Prachi & Waheed, Saman & Thakur, Shiva Kant & Sharma, Vidhi & Chanana, Arsh & Singh, Ravindra. Transdermal Drug Delivery System (Tdds) As A Novel Tool For Drug Delivery. 2023;12:688-711. DOI:10.20959/wjpr20232-26997.
 27. Wang S, Zhang D, Li C, Wang J, Zhang J, Cheng Y, Wang Q. Numerical optimization for a phase change material based lithium-ion battery thermal management system. *Applied Thermal Engineering*. 2023;222:119839.
 28. Teja SP, Damodharan N. 23 full factorial model for particle size optimization of methotrexate loaded chitosan nanocarriers: A design of experiments (DoE) approach. *BioMed research international*; 2018.
 29. Pandey Prachi Thakur Shiva Kant Chanana, Arsh Pal, Rahul. biodegradable polymer's enhancing drug delivery activity in different novel drug delivery system. *world journal of pharmacy and pharmaceutical sciences*. 2023;12:2046-2069. DOI:10.20959/wjpps2023.
 30. Fegade SL, Tande BM, Cho H, Seames WS, Sakodynskaya I, Muggli DS, Kozliak EI. Aromatization of propylene over Hzsm-5: A design of experiments (DOE) approach. *Chemical Engineering Communications*. 2013;200(8):1039-1056.
 31. Granato D, de Araújo Calado VM. The use and importance of design of experiments (DOE) in process modelling in food science and technology. *Mathematical and*

- statistical methods in food science and technology. 2014;1(1):1-18.
32. Pandey, Prachi Pal, Rahul Chanana, Arsh Singh, Ravindra. plga nanoparticle as a carrier in colon targeting drug delivery system. 2023;11:336-343. DOI:10.36673/IJRPNS.2022.v11.i05.A39.
33. Aggarwal A, Singh H, Kumar P, Singh M. Optimizing power consumption for CNC turned parts using response surface methodology and Taguchi's technique—a comparative analysis. Journal of materials processing technology. 2008;200(1-3):373-384.
34. Pandey, Prachi & Pal, Rahul & Thakur, Shiva Kant & Sharma, Vidhi & Chanana, Arsh & Singh, Ravinder. Future aspects & modification in carbon- nanoparticles in treatment and diagnosis. World Journal of Pharmacy and Pharmaceutical Sciences. 2023;12:307-324. DOI:10.20959/wjpps20232-24094.
35. Hasniyati MR, Zuhailawati H, Ramakrishnan S. A statistical prediction of multiple responses using overlaid contour plot on hydroxyapatite coated magnesium via cold spray deposition. Procedia Chemistry. 2016;19:181-188.
36. Pal, Rahul Pandey, Prachi Koli, Manju Rizwan, Mohammad & Saxena, Archita. Chitosan: As Highly Potential Biopolymer Obtainable in Several Advance Drug Delivery Systems Including Biomedical Applications "Research Paper", European Chemical Bulletin (ECB). European Chemical Bulletin. 2023;12:322-344. DOI:10.48047/ecb/2023.12.si10.0036.
37. Ho HO, Liu CH, Lin HM, Sheu MT. The development of matrix tablets for diclofenac sodium based on an empirical in vitro and *in vivo* correlation. Journal of controlled release. 1997;49(2-3): 149-156.
38. Pandey P, Pal R, Singh S, Gupta H. the current status in mucosal drug delivery system (mdds) and future prospectus in delivery: a systematic review; 2023.
39. Patra CN, Padhy S, Sen T, Bhattacharya S, Swain S, Sruti J, ME BR. Formulation development and evaluation of SR matrix tablets of stavudine. Ind J Pharm Edu Res. 2013;47(2):214.
40. Luiz MT, Viegas JSR, Abriata JP, Viegas F, de Carvalho Vicentini FTM, Bentley MVLB, Tapia-Blacido DR. Design of experiments (DoE) to develop and to optimize nanoparticles as drug delivery systems. European Journal of Pharmaceutics and Biopharmaceutics. 2021;165:127-148.
41. Pal, Rahul Pandey, Prachi Saxena, Archita Thakur, Shiva Kant. The Pharmaceutical Polymer's; A Current Status in Drug Delivery: A Comprehensive Review. Journal of Survey in Fisheries Sciences. 2023;10:3682-3692. DOI:10.53555/sfs.v10i1.1648.

© 2023 Thakur et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111654>